

INNOVATIVE DEVELOPMENTS AND RESEARCH IN EDUCATION

International scientific-online conference



CLASSIFICATION OF THE REGION OF FERGANA REGION ACCORDING TO THE LEVEL OF SALINITY

Ma'rufjonov Javohir

Student of Fergana State University

Abstract: This article provides information about salinity and the level of soil salinity in each district of Fergana region.

Key words: Saline, soil, non-saline, area, district, region, salinity

According to the description of the salinity of the soils of the Fergana region, they are divided into groups:

- 1. Not salted
- 2. Lightly salted
- 3. Moderately salted
- 4. Strongly salted
- 5. Very strongly salted

Now let's consider this level of salinity at the scale of districts. Initially, non-saline soils are 28%, slightly saline soils are 46%, moderately saline soils are 7.5%, and strongly saline soils are 18.5%. Non-saline soils are 10.8%, weakly saline soils are 50.2%, moderately saline soils are 31.8%, and strongly saline soils are 7.2% in Besharich district. In Buvaida district, non-saline soils are 12.4%, slightly saline soils are 34.8%, moderately saline soils are 33.5%, and strongly saline soils are 19.3%. Unfortunately, it is impossible to find non-saline soils in Dangara District, because all the soils in this district are saline to varying degrees. The amount of slightly saline soils is 41%, the amount of moderately saline soils is 42%, and the amount of strongly saline soils is 17%. In the Yozhiovan district of the Central Fergana region, non-saline soils are 18.3%, weakly saline soils are 52.6%, moderately saline soils are 27.7%, and strongly saline soils are 1. is 4%. Non-saline soils are 16%, slightly saline soils are 43%, moderately saline soils are 29.8%, and strongly saline soils are 11.2% in Altiariq district. In Koshtepa district, the amount of non-saline soils is 17.8%, the amount of slightly saline soils is 28.4%, the amount of moderately saline soils is 36.5%, and the amount of strongly saline soils is 17.3%. In Rishton district, non-saline soils are 17.6%, slightly saline soils are 22.1%, moderately saline soils are 12.1%, and strongly saline soils are 48.2%. If we turn our attention to the Sokh district, the amount of non-saline soils in this district is 74% and the amount of weakly absorbed soils is equal to 26%. Moderately saline and highly saline soils are not found in this area. Non-saline soils are 18.2%, slightly saline soils are 43.4%, moderately saline soils are 19.1%, and strongly saline soils are 19.3% in Taslog district. In Uchkoprik district, the amount of non-saline soils is 11.2%, the amount of weakly saline soils is 25.6%, the amount of moderately saline soils is 26.6%,



INNOVATIVE DEVELOPMENTS AND RESEARCH IN EDUCATION

International scientific-online conference



and the amount of strongly saline soils is 36.6%. Non-saline soils are 79.7%, weakly saline soils are 15%, moderately saline soils are 4.9%, strongly saline soils are 0.4% in Fergana district. In Furqat district, the amount of non-saline soils is 12.9%, the amount of weakly saline soils is 39.5%, the amount of moderately saline soils is 30.1%, and the amount of strongly saline soils is 17.5%. Non-saline soils are 49%, slightly saline soils are 27%, moderately saline soils are 22.4%, and strongly saline soils are 0.6% in the district of Uzbekistan. In Kuva District, non-saline soils are 34.2%, slightly saline soils are 56.5%, moderately saline soils are 9.2%, and strongly saline soils are only 0.1%.

If we consider these indicators by region, the amount of non-saline soils is 27.9%, the amount of weakly saline soils is 35.6%, the amount of moderately saline soils is 22.2%, strongly the amount of saline soils is 14.3%. Fergana District is the region with the most widespread non-saline soils, 79.7%. The region with weakly saline soils is the most widespread with 56.5% in Kuva district. Dangara district is the clear leader in the region where moderately saline soils are most common with 42%. It can be seen that Rishton district is in the first place among regions with strong salinity with 48.2%. The least common area of highly saline soils is Kuva district with 0.1%, and as mentioned above, strongly saline soils are hardly found in Dangara district.

REFERENCES:

- 1. Iminchayev R.A Joʻrayeva M.M, Ismoilov M.I, Ma'rufjonov J.Gʻ Fargʻona vodiysi sharoitida "Polovchanka" bugʻdoy navini oziqlanish tartibotining iqtisodiy samaradorligi «Science and innovation»
- 2. Sotiboldiyeva G, Ma'rufjonov J, Solijonova D, Toshpo'latova Y. Kaliliy o'g'it konlari va uning ahamiyati. "Modern Science and Scientific Studies" 91-93 b
- 3. Sotiboldieva, G., Abduxakimova, X., Mirzakarimova, I., Xojiboev, B., & Qirgizova, M. (2022). Sugʻoriladigan boʻz tuproqlar mintaqasida kalsiyning biogeokimyoviy xususiyatlari. Science and innovation, 1(A7), 121-126.
- 4. Yuldashev, G., & Sotiboldieva, G. (2015). Formation of the absorbed foundations of the irrigated gray-brown soils of the Sokhsky cone of carrying out. Europaische Fachhochschule, (5), 3-6.
- 5. Sotiboldieva, G., Abduxakimova, X., Yuldashev, A., & Xasanov, R. (2022). Sugʻoriladigan kolmatajlangan boʻz tuproqlarda stronsiyning pedogeokimyosi. Science and innovation, 1(D7), 140-145.
- 6. Yuldashev, G., & Sotiboldieva, G. (2015). Formation of the absorbed foundations of the irrigated gray-brown soils of the Sokhsky cone of carrying out. Europaische Fachhochschule, (5), 3-6.
- 7. Юлдашев F, C. Г. (2015). Кольматажланган тупрокларда стронций ва барий. УзМУ хабарлари, 3(2), 138-143.



INNOVATIVE DEVELOPMENTS AND RESEARCH IN EDUCATION International scientific-online conference



- 8. Сотиболдиева, Г., & Абдуллаева, Л. (2020). Сух ва Исфайрамсой дарё ёйилмаларида шаклланган сугориладиган кольматажланган тупрокларнинг галогенетик хусусиятларини тавсифи. Илм-фан ва таълимнинг ривожланиш истикболлари мавзусидаги илмий канфренция туплами. www. openscience. uz, 27, 309-313.
- 9. Юлдашев, Г., Исагалиев, М., Сотиболдиева, Г., & Турдалиев, А. БИОМИКРОЭЛЕМЕНТЫ В АГРОЛАНДШАФТАХ ЦЕНТРАЛЬНОЙ ФЕРГАНЫ. СЕМИНАР—КРУГЛЫЙ СТОЛ 6. ПРИЁМЫ РЕГУЛИРОВАНИЯ ПОЧВЕННОГО ПЛОДОРОДИЯ И ОХРАНА ПОЧВЕННЫХ РЕСУРСОВ, 409.
- 10. Toshmirzayeva, G., & Sotiboldiyeva, G. (2021, July). LIGHT GRAY AND TYPICAL GRAY SOILS OF UCHKURGAN DISTRICT. In Конференции.
- 11. Sotiboldiyeva, G. T. (2018). Farg 'ona viloyati kolmatajlangan tuproqlarining biogeokimyoviy xususiyatlari va ulardan foydalanish. Diss. bffd-Toshkent, 31-42.
- 12. Yuldashev, G., & Sotiboldiyeva, G. (2021). BIOGEOCHEMISTRY OF SELENIUM AND ARSENIC IN AGRICULTURAL LANDSCAPES: https://doi.org/10.47100/conferences. v1i1. 1363. In RESEARCH SUPPORT CENTER CONFERENCES (No. 18.06).
- 13. Sotiboldiyeva, G., Abdukhakimova, K., & Niyozov, Q. (2021). About digital mapping of biomicroelements: https://doi. org/10.47100/conferences. v1i1. 1366. In RESEARCH SUPPORT CENTER CONFERENCES (No. 18.06).
- 14. Сотиболдиева, Г. Т. (2018). Фаргона вилояти кольматажланган тупрокларининг биогеокимёвий хусусиятлари ва улардан фойдаланиш.: дисс. Автореф. б. ф. ф. д.(PhD)-Т.
- 15. Sotiboldieva, G. T., & Yuldashev, G. Y. (2014). POLLUTION OF IRRIGATED SOILS IN THE SEROZEM ZONE BY RADIONUCLIDES. The Way of Science, 33.
- 16. Турдалиев, А., & Сотиболдиева, Г. Агрохимические свойства трудномелиорируемых почв Ферганы.
- 17. Юлдашев, Г., Холдарова, М., Исагалиев, М., Турдалиев, А., & Сотиболдиева, Г. (2013). Агрохимические свойства трудномелиорируемых почв Ферганы. Аграрный вестник Урала, (3 (109)), 16-17.
- 18. Юлдашев, Г., Исагалиев, М., Аскаров, Х., & Сотиболдиева, Г. (2016). Агрофизические свойства бурых горно-лесных почв Западной Ферганы. Почвоведение-продовольственной и экологической безопасности страны, 397-398.
- 19. Sotiboldiyeva Goʻzalxon Tolibjonovna. TUPROQSHUNOSLIK YOʻNALISHI TALABALARI UCHUN MALAKAVIY AMALIYOTINI TASHKILLASH METODIKASI. World of Science. 2023/4/20. 142-14
- 20. Sotiboldiyeva, G. (2023). KOLMATAJLANGAN SUR TUSLI QO 'NGIR TUPROQ VA GRUNTLARNING MEXANIK TARKIBI. Science and innovation, 2(Special Issue 6), 834-838.